

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/17, 39/395 // C07K 16/28, 14/705	A2	(11) International Publication Number: WO 00/15247 (43) International Publication Date: 23 March 2000 (23.03.00)
(21) International Application Number: PCT/US99/21170 (22) International Filing Date: 13 September 1999 (13.09.99) (30) Priority Data: 60/100,182 14 September 1998 (14.09.98) US (71) Applicant (for all designated States except US): BIOGEN, INC. [US/US]; 14 Cambridge Center, Cambridge, MA 02142 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MUNDY, Gregory, R. [US/US]; 3719 Morgan's Creek, San Antonio, TX 78230 (US). YONEDA, Toshiyuki [JP/US]; 3530 Hunters Sound, San Antonio, TX 78230 (US). (74) Agent: KAPLAN, Warren, A.; Biogen, Inc., 14 Cambridge Center, Cambridge, MA 02142 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHODS OF TREATING MULTIPLE MYELOMA AND MYELOMA-INDUCED BONE RESORPTION USING INTEGRIN ANTAGONISTS (57) Abstract Antagonists of alpha4 integrin/alpha4 integrin ligand adhesion, which inhibit the biological effects of such adhesion are described and methods for their use are detailed. Such antagonists are useful in suppressing bone destruction associated with multiple myeloma. The homing of multiple myeloma cells to bone marrow and their alpha4 integrin-dependent release of bone-resorbing factors, resulting in bone destruction in patients with multiple myeloma, is inhibited.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 **Methods of Treating Multiple Myeloma and Myeloma-induced Bone Resorption Using Integrin Antagonists**

FIELD OF THE INVENTION

10 The present invention relates to a treatment for multiple myeloma, and the release of bone-resorbing factors by myeloma cells, resulting in severe bone loss, which is the major side-effect of myeloma in man. More particularly, this invention relates to integrin antagonists, such as antagonists of alpha4 containing integrins, which inhibit the biological effects of such adhesion, associated with homing of multiple myeloma
15 cells to bone marrow; their subsequent integrin-dependent survival; and their integrin-dependent release of bone-resorbing factors, resulting in bone destruction in patients with multiple myeloma.

20 **BACKGROUND OF THE INVENTION**

 Multiple myeloma is the second most common hematologic malignancy, with 15,000 new cases diagnosed each year and 30,000 to 40,000 myeloma patients in the U.S. annually (Mundy and Bertolini 1986). Eighty percent of the patients suffer from
25 devastating osteolytic bone destruction caused by increased osteoclast (OCL) formation and activity (Mundy and Bertolini 1986). This bone destruction can cause excruciating bone pain, pathologic fractures, spinal cord compression, and life-threatening hypercalcemia. Because multiple myeloma cannot be cured by standard chemotherapy or stem cell transplantation (Attal et al, 1996), and because of the severe morbidity and
30 potential mortality associated with myeloma bone disease, treatment strategies that control the myeloma growth itself, and in particular the osteolytic bone destruction that occurs in these patients, are vitally important.

 However, the pathologic mechanisms responsible for the increased osteoclast activity in patients with multiple myeloma are unknown (Mundy, 1998). The bone
35 lesions occur in several patterns. Occasionally, patients develop discrete osteolytic

5 lesions that are associated with solitary plasmacytomas. Some patients have diffuse osteopenia, which mimics the appearance of osteoporosis, and is due to the myeloma cells being spread diffusely throughout the axial skeleton. In most patients there are multiple discrete lytic lesions occurring adjacent to nests of myeloma cells.

Hypercalcemia occurs as a consequence of bone destruction in about one-third of
10 patients with advanced disease. Rarely, patients with myeloma do not have lytic lesions or bone loss, but rather have an increase in the formation of new bone around myeloma cells. This rare situation is known as osteosclerotic myeloma.

Osteolytic bone lesions are by far the most common skeletal manifestations in patients with myeloma (Mundy, 1998). Although the precise molecular mechanisms
15 remain unclear, observations over 15 years have shown that: 1) The mechanism by which bone is destroyed in myeloma is via the osteoclast, the normal bone-resorbing cell; 2) Osteoclasts accumulate on bone-resorbing surfaces in myeloma adjacent to collections of myeloma cells and it appears that the mechanism by which osteoclasts are stimulated in myeloma is a local one; 3) It has been known for many years that cultures
20 of human myeloma cells in vitro produce several osteoclast activating factors, including lymphotoxin-alpha (LT-a), interleukin-1 (IL-1), parathyroid-hormone related protein (PTHrP) and interleukin-6 (IL-6); 4) Hypercalcemia occurs in approximately one-third of patients with myeloma some time during the course of the disease. Hypercalcemia is always associated with markedly increased bone resorption and frequently with
25 impairment in glomerular filtration; 5) The increase in osteoclastic bone resorption in myeloma is usually associated with a marked impairment in osteoblast function. Alkaline phosphatase activity in the serum is decreased or in the normal range, unlike patients with other types of osteolytic bone disease, and radionuclide scans do not show evidence of increased uptake, indicating impaired osteoblast responses to the increase in
30 bone resorption.

Although various mediators listed above have been implicated in the stimulation of osteoclast activity in patients with multiple myeloma, reports of factors produced by myeloma cells have not been consistent, and some studies have been inconclusive due to the presence of other contaminating cell types, including stromal cells and
35 macrophages, in the multiple myeloma cell population. IL-6 is a major myeloma growth

5 factor that enhances the growth of several myeloma cell lines and freshly isolated
myeloma cells from patients (Bataille et al., 1989). IL-6 production can be detected in
about 40% of freshly isolated myeloma cells by PCR, but only 1 in 150 patients studied
show detectable IL-6 production by immunocytochemistry or ELISA assays (Epstein
1992). The IL-6 receptors were only detected in 6 of 13 samples from patients with
10 multiple myeloma (Bataille et al, 1992). Furthermore, mature myeloma cells have been
reported to have a minimal proliferative response to IL-6. Interleukin-11 (IL-11) has an
IL-6-like activity on plasmacytomas, but to date no one has demonstrated that myeloma
cells produce IL-11. Bataille and coworkers (1995) have shown that perfusion of 5
patients with refractory myeloma with an antibody to IL-6 decreased the size of the
15 myeloma cell burden in only 2 of these patients. IL-1 is an extremely potent bone-
resorbing agent that induces hypercalcemia in animal models in the absence of renal
failure (Boyce et al, 1989). In contrast, hypercalcemia rarely occurs in myeloma
patients without renal failure. More importantly, in highly purified myeloma cells, no
IL-1 and only rare TNF-a production can be detected, suggesting that other
20 contaminating cell types such as macrophages may be the source of IL-1 and TNF-a
(Epstein 1992). Similarly, LT-a is produced by most human myeloma cell lines
(Bataille et al, 1995) but does not appear to be produced by myeloma cells in vivo
(Alsina et al, 1996). In addition to IL-1, TNF-a, LT-a, and IL-6, myeloma cells produce
a truncated form of M-CSF which is biologically active, but M-CSF does not cause
25 hypercalcemia or induce osteoclast formation by itself in human marrow cultures
(MacDonald et al, 1986).

Thus, the role of any of these factors in osteolytic bone disease in patients with
myeloma has not been clearly demonstrated in vivo, so that known cytokines clearly do
not totally account for the bone resorption seen in these patients.

Role of Adhesive Molecule Interactions in Myeloma Bone Disease

5 Anderson and coworkers were the first group to demonstrate the importance of
adhesive interactions between myeloma cells and cells in the marrow microenvironment
both in the growth of myeloma cells and the development of osteolytic bone disease.
Multiple myeloma cells express cell surface adhesion molecules, CD29 (VLA-4), LFA-
1, and CD44 (Chauhan et al, 1995). These workers suggested that myeloma cells
10 localized to the marrow via specific adhesion interactions between extracellular matrix
proteins and bone marrow stromal cells. They further showed that adhesion of multiple
myeloma cells to stromal cells triggered IL-6 secretion by both normal and multiple
myeloma bone marrow-derived stromal cells and increased IL-6-mediated tumor cell
growth. However, antibodies to CD29, LFA-1 or CD44 did not decrease IL-6
15 production by marrow stromal cells in response to myeloma cells, suggesting that
another ligand-receptor interaction triggered the IL-6 secretion by bone marrow stromal
cells binding to myeloma cells. Mere identification of a possible adhesion pathway does
not necessarily mean that the pathway is important. In this case none of the implicated
pathways plays a role in IL-6 production.

20 Vanderkerken et al (1997) also examined the phenotypic adhesion profile of
murine 5T2 cells and 5T33 myeloma cells in a model of murine myeloma. These
investigators showed that these cell lines expressed VLA-4, VLA-5, LFA-1, and CD44,
and suggested that these adhesive interactions might be important for myeloma cells to
bind to marrow stromal cells.

25 Nevertheless, despite many laboratory advances, the fundamental mechanisms
underlying increased osteoclastic bone destruction in myeloma in vivo remain poorly
understood. This is reflected in the inability to easily translate the data on adhesive
interactions obtained in vitro to the in vivo setting. For example, many in vitro studies
implicate both the integrin VLA-4 and the integrin LFA-1 in the adhesion of
30 hematopoietic stem cells to bone marrow stroma (reviewed in Papayannopoulou and
Nakamoto, 1993). These in vitro data would predict that either pathway, if blocked in
vivo, would result in peripheralization of hematopoietic stem cells from marrow to
peripheral blood. Yet, in a primate study, while a monoclonal antibody (mAb) to VLA-
4 effectively peripheralized stem cells, a monoclonal antibody to the beta2 integrin
35 chain of LFA-1 was without effect, despite increasing neutrophil counts, thus

5 demonstrating the efficacy of the mAb (Papayannopoulou and Nakamoto, 1993). These data show that the in vitro results were in fact unable to accurately predict in vivo relevance.

It should be noted that the role of integrin VLA-4 has been studied in metastasis of multiple tumors, including leukemias such as lymphoma, with contradictory results. Thus, transfection of the human alpha 4 chain into Chinese Hamster Ovary (CHO) cells resulted in VLA-4 expression, and rendered these cells able to migrate to bone marrow in vivo, a phenomenon inhibited by mAbs to VLA-4 (Matsuura et al, 1996). In contrast, transfection of lymphoma cells with VLA-4 strongly inhibited metastasis to liver, lung and kidney, and was without effect on homing and proliferation in marrow (Gossler et al., 1996). In addition, expression of VLA-4 on highly metastatic murine melanoma cells strongly inhibited the formation of pulmonary metastases in vivo (Qian et al., 1994), and did not predispose melanoma to bone marrow metastasis.

In summary it is not clear on the basis of in vitro studies, how to reliably predict in vivo relevance of adhesion pathways. Furthermore, even when in vivo studies have been performed, the resultant data are inconsistent. One major reason for the perplexing inconsistencies in the field of multiple myeloma is that currently available animal models are not good predictors of human disease. In the case of multiple myeloma, human and murine myeloma cell lines which can be grown in vitro rarely are associated with bone destruction in vivo (Mundy 1998).

25 It would be highly desirable to identify compounds or antagonists which inhibit production of these bone-resorbing factors, thus halting progressive bone destruction and improving the quality of life of patients with myeloma.

SUMMARY OF THE INVENTION

30 We have used a recently developed murine model of multiple myeloma in which the mouse develops severe osteolysis with all the hallmarks of human disease (Garrett 1997). Using this cell line and animal model we have established that inhibition of the alpha4 integrin/alpha4 integrin ligand pathway in vivo leads to reduced capacity for multiple myeloma cells to proliferate and/or survive. We show that cell-cell attachment between myeloma cells and marrow stromal cells via the VLA-4/VCAM-1 interaction

5 is required for an increase in the production of an activity which stimulates osteoclastic bone resorption in the bone microenvironment in vitro.

We propose that this interaction is critical to the homing of myeloma cells to the marrow compartment, to their subsequent survival and growth, to ultimately to the progression of myeloma-induced osteolysis. We tested this in the animal model and found that, in vivo, an antagonist of the alpha4 subunit-containing integrin VLA-4 –
10 strongly inhibits the production of antibody of the IgG2b subtype. This isotype is the same as that produced by the 5TGM1 cell line, and is an accurate surrogate for the number of myeloma cells in the marrow compartment at any time. Thus, blockade of the VLA-4 pathway strongly inhibits IgG2b production, and by implication, the level of myeloma burden.
15

One aspect of the invention is a method for the treating multiple myeloma comprising administering to an individual a therapeutically effective amount of a composition comprising an antagonist of an interaction between an integrin with an alpha4 subunit (e.g., VLA-4) and a ligand for this integrin (e.g., VCAM-1). This
20 antagonist can be an alpha4 integrin binding agent or an alpha4 integrin ligand binding agent. Preferred agents are anti-VLA4 or anti-alpha4beta7 antibody homologs (human antibody, a chimeric antibody, a humanized antibody and fragments thereof); anti-VCAM-1 antibody homologs (a human antibody, a chimeric antibody, a humanized antibody and fragments thereof); and a small molecule inhibitor of interactions of
25 alpha4 subunit containing integrins with their ligands. The composition can be administered at a dosage so as to provide from about 0.1 to about 20 mg/kg body weight. In particular, the preferred agents can antagonize an interaction: a) of both VLA-4 and alpha4 beta 7 collectively with their respective alpha4 ligands; or b) only of VLA-4 with its alpha4 ligand; or c) only of alpha4beta7 with its alpha4 ligand.
30

Another aspect of the invention is a method for inhibiting bone resorption associated with tumors of bone marrow, the method comprising administering to a mammal with said tumors an antagonist of an interaction between an alpha4 subunit-containing integrin such as VLA-4 and a ligand for this alpha4 subunit containing
35 integrin, such as VCAM-1, in an amount effective to provide inhibition of the bone

5 resorption. . This antagonist can be an alpha4 integrin binding agent such as a VLA-4
binding agent or an alpha4 integrin ligand binding agent such as a VCAM-1 binding
agent. Preferred agents are anti-VLA4 or anti alpha4beta7 antibody homologs (human
antibody, a chimeric antibody, a humanized antibody and fragments thereof); anti-
VCAM-1 antibody homologs (a human antibody, a chimeric antibody, a humanized
10 antibody and fragments thereof); and a small molecule inhibitor of the interaction of
alpha4 subunit-containing integrins with their respective alpha4 integrin ligands (e.g,
the VCAM-1/VLA-4 interaction). The antagonist can be administered at a dosage so as
to provide from about 0.1 to about 20 mg/kg body weight.

15 Yet another aspect of the invention is a method of treating a subject having a
disorder characterized by the presence of osteoclastogenesis, the method comprising
administering to the subject an antagonist of an interaction between an alpha4 subunit-
bearing integrin and a ligand for an alpha4 subunit-bearing integrin, in an amount
sufficient to suppress the osteoclastogenesis. Similarly, the antagonist can be a alpha4
binding agent or an alpha4 ligand binding agent. Preferred agents are anti-VLA4 or
20 anti-alpha4beta7 antibody homologs (human antibody, a chimeric antibody, a
humanized antibody and fragments thereof); anti-VCAM-1 antibody homologs (a
human antibody, a chimeric antibody, a humanized antibody and fragments thereof);
and a small molecule inhibitor of the interaction of alpha4 subunit-containing integrins
with their respective alpha4 integrin ligands (e.g, the VCAM-1/VLA-4 interaction).
25 The composition can be administered at a dosage so as to provide from about 0.1 to
about 20 mg/kg body weight. Unless stipulated otherwise, all references are
incorporated herein by reference.

30 BRIEF DESCRIPTION OF THE FIGURES

Figure 1.

**Effect of Neutralizing Antibodies on TRAP-positive Multinucleated OC-like Cell
Formation in the Co-cultures of 5TGM1 cells and Bone Marrow Cells.**

5 A mixture of 5TGM1 cells (1×10^3) and marrow cells (1×10^6) in suspension was
plated in 48-well culture plates and cultured with or without 10 ug/ml anti-VCAM-1
antibody (VCAM-1 Ab), anti- $\alpha 4 \beta 1$ antibody ($\alpha 4 \beta 1$ Ab), anti-ICAM-1 antibody
(ICAM-1 Ab) or rat IgG as a control. After 6 days of culture, cultures were fixed and
the number of TRAP-positive multinucleated OC-like cells (TRAP(+) MNC)
10 determined. Both VCAM-1 Ab and $\alpha 4 \beta 1$ Ab inhibited TRAP(+) MNC
formation, while ICAM-1 Ab had no effect. Data are expressed as mean \pm S.E. (n=3). *
= Significantly different from IgG control.

15

Figure 2

**Effect of 5TGM1 and ST2 Conditioned Media on bone resorption in Organ
20 Cultures of Fetal Rat Long Bones.**

Conditioned media (48 hours) obtained from ST2 alone, 5TGM1 alone, and co-
cultures of ST2 and 5TGM1 were assayed for bone resorbing activity in organ cultures
of 45 calcium-labeled fetal rat long bones. Labeled fetal rat long bones were cultured in
the presence of conditioned media (40%v/v) or control medium for 120 hours. Data are
25 expressed as percentage increase of calcium release over than in the control medium.
Release from conditioned medium of ST2 stromal cells is shown as the open bar.
Release from 5TGM1 is the hatched bar. Release from conditioned medium harvested
from co-culture of 5TGM1 and ST2 is the closed bar. Data are expressed as mean \pm
S.E. (n=4). * = significantly different from ST2 alone. *** = significantly different
30 from 5TGM1 alone.

Figure 3

**Effect of Recombinant Soluble VCAM-1 (sVCAM-1) on the Production of
Osteoclastogenic Activity by 5TGM1 Cells.**

5 Conditioned medium was harvested from 5TGM1 cells cultured in the presence
or absence of sVCAM-1 (1×10^{-8} to 1×10^{-7} Molar) for 24 hours. Osteoclastogenic
activity of these conditioned media was assayed in the mouse marrow cultures. Bone
marrow cells (1×10^6 /well) were plated into 48-well plates, and cultured in the presence of
10 conditioned media (hatched bars) or control medium (IMDM) containing the same
concentrations of sVCAM-1 (open bars). After 6 days, cultures were fixed and the —
number of TRAP-positive multinucleated OC-like cells (TRAP+ MNC) was
determined. Conditioned medium from 5TGM1 cells treated with 1×10^{-7} M sVCAM-1
increased TRAP(+)MNC formation. Data are expressed as mean \pm S.E. (n=3). * =
significantly different from controls.

15 **Figure 4**

Effect of mAb PS2 to VLA-4 on serum IgG2b elevation in 5TGM1-bearing mice

Mice were injected with 1×10^5 5TGM1 cells, which were allowed to colonize the bone
20 marrow. Mice were split into two groups of three, one serving as a control group, and
the second treated on days 8, 11, 14, 17, and 20 with 80 ug mAb PS/2 (~4 mg/kg).
Levels of IgG2b, the antibody isotype produced by 5TGM1 myeloma cells, were
measured weekly from weeks 1 to 6. Mab treatment strongly inhibited IgG2b
production, indicative of inhibition of myeloma cell survival and growth in vivo.

25 **Figure 5**

30 **Effect of mAb M/K-2.7 to VCAM-1 on serum IgG2b elevation in 5TGM1-bearing
mice**

Mice were injected with 5TGM1 cells as described in Figure 4, which were allowed to
colonize the bone marrow. Mice were split into groups of four or five, one group
35 serving as a control group (open square), the second/third groups treated

5 prophylactically at 80 ug (open diamonds) and 160 ug mAb (open circles) (~4 to 8 mg/kg), the fourth group treated therapeutically at 160 ug mAb (triangles). Levels of IgG2b, the antibody isotype produced by 5TGM1 myeloma cells, were measured. Mab treatment strongly inhibited IgG2b production, indicative of inhibition of myeloma cell survival and growth in vivo.

10 **Figure 6 Effect of anti-alpha4 Integrin Antibody on Survival of Multiple Myeloma-bearing Mice**

DETAILED DESCRIPTION OF THE INVENTION

15 The invention relates to treatments for, among other things, preventing multiple myeloma. More particularly, methods of the invention relate to the use of antagonists of an interaction between an integrin containing an alpha4 subunit and a ligand for this integrin in the treatment of multiple myeloma. The term "multiple myeloma" is intended to mean a medical condition in an individual having a neoplastic disease of plasma cells, with the neoplastic clone representing cells at different stages in the
20 plasma cell lineage from patient to patient (Mundy, 1998).

Alpha 4 beta 1 integrin is a cell-surface receptor for VCAM-1, fibronectin and possibly other molecules that bind with, or otherwise interact with, alpha 4 beta 1 integrin. In this regard, such molecules that bind with, or otherwise interact with, alpha
25 4 subunit containing integrin are individually and collectively referred to as "alpha4 ligand(s)". The term alpha4 beta 1 integrin ("VLA-4" or "alpha4 beta 1" or "alpha4 beta 1 integrin", used interchangeably) herein thus refers to polypeptides which are capable of binding to VCAM-1 and members of the extracellular matrix proteins, most particularly fibronectin, or homologs or fragments thereof, although it will be appreciated by
30 workers of ordinary skill in the art that other ligands for VLA-4 may exist and can be analyzed using conventional methods.

Nevertheless, it is known that the alpha4 subunit will associate with other beta subunits besides beta1 so that we may define the term "alpha 4 integrin" as being those integrins whose alpha4 subunit associates with one or another of the beta subunits. A
35 further example of an "alpha4" integrin is alpha4beta7 (R. Lobb and M. Hemler, 1994).

5 As used herein, the term "alpha4 integrin(s)" means VLA-4, as well as integrins that contain beta 1, beta7 or any other beta subunit.

As discussed herein, the antagonists used in methods of the invention are not limited to a particular type or structure of molecule so that, for purposes of the invention, any agent capable of binding to any integrin containing an alpha 4 subunit
10 such as VLA-4 on the surface of VLA-4 bearing cells and/or alpha4beta7 integrin on the surface of alpha4beta7-bearing cells [see Lobb and Hemler, J. Clin. Invest., 94: 1722-1728 (1994)] and/or to their respective alpha4 ligands such as VCAM-1 and MadCAM, respectively, on the surface of VCAM-1 and MadCAM bearing cells, and which effectively blocks or coats VLA-4 (or alpha4beta7) or VCAM-1 (or MadCAM)
15 (i.e., a "an alpha4 integrin binding agent" and "alpha4 integrin ligand binding agent" respectively), is considered to be an equivalent of the antagonists used in the examples herein.

An integrin "antagonist" includes any compound that inhibits an alpha 4 integrin(s) from binding with an alpha 4 integrin ligand and/or receptor. Anti-integrin antibody or antibody homolog-containing proteins (discussed below) as well as other
20 molecules such as soluble forms of the ligand proteins for integrins are useful. Soluble forms of the ligand proteins for alpha4 integrins include soluble VCAM-1 or collagen peptides, VCAM-1 fusion proteins, or bifunctional VCAM-1/Ig fusion proteins. For example, a soluble form of an alpha4 integrin ligand or a fragment thereof may be
25 administered to bind to integrin, and preferably compete for an integrin binding site on cells, thereby leading to effects similar to the administration of antagonists such as anti-alpha4 integrin (e.g., alpha4 beta7 antibodies and/or VLA-4 antibodies. In particular, soluble alpha4 integrin mutants that bind alpha 4 integrin ligand but do not elicit integrin-dependent signaling are included within the scope of the invention. Such
30 mutants can act as competitive inhibitors of wild type integrin protein and are considered "antagonists". Other antagonists used in the methods of the invention are "small molecules", as defined below.

Included within the invention are methods using an agent that antagonizes the action of more than one alpha 4 integrin, such as a single small molecule or antibody
35 homolog that antagonizes several alpha 4 integrins such as VLA-4 and alpha4 beta 7, or

5 other combinations of alpha4 integrins. Also included within the scope of the invention are methods using a combination of different molecules such that the combined activity antagonizes the action of more than one alpha4 integrin, such as methods using several small molecules or antibody homologs that in combination antagonize the alpha 4 integrins VLA-4 and alpha4 beta 7, or other combinations of integrins.

10 As discussed herein, certain integrin antagonists can be fused or otherwise — conjugated to, for instance, an antibody homolog such as an immunoglobulin or fragment thereof and are not limited to a particular type or structure of an integrin or ligand or other molecule. Thus, for purposes of the invention, any agent capable of forming a fusion protein (as defined below) and capable of binding to alpha4 integrin
15 ligands and which effectively blocks or coats alpha4 beta 7 and/or VLA-4 integrin is considered to be an equivalent of the antagonists used in the examples herein.

For the purposes of the invention an “antagonist of the alpha 4 integrin ligand/alpha4 integrin interaction” refers to an agent, e.g., a polypeptide or other molecule, which can inhibit or block alpha 4 ligand (e.g., VCAM-1) and/or alpha 4
20 integrin (e.g., alpha4beta7 or VLA-4) -mediated binding or which can otherwise modulate alpha4 ligand and/or alpha4 integrin function, e.g., by inhibiting or blocking alpha4-ligand mediated alpha4 integrin signal transduction or alpha4 ligand- mediated alpha4 ligand signal transduction and which is effective in the treatment of multiple myeloma, preferably in the same manner as are anti-alpha4 integrin antibodies.

25 Specifically, an antagonist of the VCAM-1/ VLA-4 interaction is an agent which has one or more of the following properties: (1) it coats, or binds to, VLA-4 on the surface of a VLA-4 bearing cell (e.g., a myeloma cell) with sufficient specificity to inhibit a VLA-4-ligand/VLA-4 interaction, e.g., the VCAM-1/VLA-4 interaction between bone stromal cells and myeloma cells; (2) it coats, or binds to, VLA-4 on the
30 surface of a VLA-4 bearing cell (i.e., a myeloma cell) with sufficient specificity to modify, and preferably to inhibit, transduction of a VLA-4-mediated signal e.g., VLA-4/VCAM-1-mediated signaling; (3) it coats, or binds to, a VLA-4-ligand, (e.g., VCAM-1) on bone stromal cells with sufficient specificity to inhibit the VLA- 4 /VCAM interaction; (4) it coats, or binds to, a VLA-4-ligand (e.g., VCAM- 1) on bone stromal
35 cells with sufficient specificity to modify, and preferably to inhibit, transduction of

5 VLA-4-ligand mediated VLA-4 signaling, e.g., VCAM-1-mediated VLA-4 signaling. In preferred embodiments the antagonist has one or both of properties 1 and 2. In other preferred embodiments the antagonist has one or both of properties 3 and 4. Moreover, more than one antagonist can be administered to a patient, e.g., an agent which binds to VLA-4 can be combined with an agent which binds to VCAM-1.

10 For example, antibodies or antibody homologs (discussed below) as well as — soluble forms of the natural binding proteins for VLA-4 and VCAM-1 are useful. Soluble forms of the natural binding proteins for VLA-4 include soluble VCAM-1 peptides, VCAM-1 fusion proteins, bifunctional VCAM-1/Ig fusion proteins, fibronectin, fibronectin having an alternatively spliced non-type III connecting segment,
15 and fibronectin peptides containing the amino acid sequence EILDV or a similar conservatively substituted amino acid sequence. Soluble forms of the natural binding proteins for VCAM-1 include soluble VLA-4 peptides, VLA-4 fusion proteins, bifunctional VLA-4/Ig fusion proteins and the like. As used herein, a “soluble VLA-4 peptide” or a “soluble VCAM-1 peptide” is an VLA-4 or VCAM-1 polypeptide
20 incapable of anchoring itself in a membrane. Such soluble polypeptides include, for example, VLA-4 and VCAM polypeptides that lack a sufficient portion of their membrane spanning domain to anchor the polypeptide or are modified such that the membrane spanning domain is non-functional. These binding agents can act by competing with the cell-surface binding protein for VLA-4 or by otherwise altering
25 VLA-4 function. For example, a soluble form of VCAM-1 (see, e.g., Osborn et al. 1989, Cell, 59: 1203-1211) or a fragment thereof may be administered to bind to VLA-4, and preferably compete for a VLA-4 binding site on myeloma cells, thereby leading to effects similar to the administration of antagonists such as small molecules or anti-VLA-4 antibodies.

30 In another example, VCAM-1, or a fragment thereof which is capable of binding to VLA-4 on the surface of VLA-4 bearing myeloma cells, e.g., a fragment containing the two N-terminal domains of VCAM-1, can be fused to a second peptide, e.g., a peptide which increases the solubility or the in vivo life time of the VCAM-1 moiety. The second peptide can be a fragment of a soluble peptide, preferably a human peptide,
35 more preferably a plasma protein, or a member of the immunoglobulin superfamily. In

5 particularly preferred embodiments the second peptide is IgG or a portion or fragment thereof, e.g., the human IgG1 heavy chain constant region and includes, at least the hinge, CH2 and CH3 domains.

Other antagonists useful in the methods of the invention include, but are not limited to, agents that mimic the action of peptides (organic molecules called "small
10 molecules") capable of disrupting the $\alpha 4$ integrin/ $\alpha 4$ integrin ligand interaction by, for instance, blocking VLA-4 by binding VLA-4 receptors on the surface of cells or blocking VCAM-1 by binding VCAM-1 receptors on the surface of cells. These "small molecules" may themselves be small peptides, or larger peptide-containing organic compounds or non-peptidic organic compounds. A "small molecule", as defined herein,
15 is not intended to encompass an antibody or antibody homolog. Although the molecular weight of such "small molecules" is generally less than 2000, we don't intend to apply this figure as an absolute upper limit on molecular weight.

For instance, small molecules such as oligosaccharides that mimic the binding domain of a VLA-4 ligand and fit the receptor domain of VLA-4 may be employed.
20 (See, J.J. Devlin et al., 1990, Science 249: 400-406 (1990), J.K. Scott and G.P. Smith, 1990, Science 249: 386-390, and U.S. Patent 4,833,092 (Geysen), all incorporated herein by reference. Conversely, small molecules that mimic the binding domain of a VCAM-1 ligand and fit the receptor domain of VCAM-1 may be employed.

Examples of other small molecules useful in the invention can be found in
25 Komoriya et al. ("The Minimal Essential Sequence for a Major Cell Type-Specific Adhesion Site (CS1) Within the Alternatively Spliced Type III Connecting Segment Domain of Fibronectin Is Leucine-Aspartic Acid-Valine", J. Biol. Chem., 266 (23), pp. 15075-79 (1991)). They identified the minimum active amino acid sequence necessary to bind VLA-4 and synthesized a variety of overlapping peptides based on the
30 amino acid sequence of the CS-1 region (the VLA-4 binding domain) of a particular species of fibronectin. They identified an 8-amino acid peptide, Glu-Ile-Leu-Asp-Val-Pro-Ser-Thr, as well as two smaller overlapping pentapeptides, Glu-Ile-Leu-Asp-Val and Leu-Asp-Val-Pro-Ser, that possessed inhibitory activity against fibronectin-dependent cell adhesion. Certain larger peptides containing the LDV sequence were
35 subsequently shown to be active in vivo (T. A. Ferguson et al., "Two Integrin Binding

5 Peptides Abrogate T-cell-Mediated Immune Responses In Vivo", Proc. Natl. Acad. Sci. USA, 88, pp. 8072-76 (1991); and S. M. Wahl et al., "Synthetic Fibronectin Peptides Suppress Arthritis in Rats by Interrupting Leukocyte Adhesion and Recruitment", J. Clin. Invest., 94, pp. 655-62 (1994)). A cyclic pentapeptide, Arg-Cys-Asp-TPro-Cys (wherein TPro denotes 4-thioproline), which can inhibit both VLA-4 and VLA-5
10 adhesion to fibronectin has also been described. (See, e.g., D.M. Nowlin et al. "A Novel Cyclic Pentapeptide Inhibits Alpha4Beta1 Integrin-mediated Cell Adhesion", J. Biol. Chem., 268(27), pp. 20352-59 (1993); and PCT publication PCT/US91/04862). This pentapeptide was based on the tripeptide sequence Arg-Gly-Asp from FN which had been known as a common motif in the recognition site for several extracellular-matrix
15 proteins.

 Examples of other small molecule VLA-4 inhibitors have been reported, for example, in Adams et al. "Cell Adhesion Inhibitors", PCT US97/13013, describing linear peptidyl compounds containing beta-amino acids which have cell adhesion inhibitory activity. International patent applications WO 94/15958 and WO 92/00995
20 describe cyclic peptide and peptidomimetic compounds with cell adhesion inhibitory activity. International patent applications WO 93/08823 and WO 92/08464 describe guanidiny-, urea- and thiourea-containing cell adhesion inhibitory compounds. United States Patent No. 5,260,277 describes guanidiny cell adhesion modulation compounds.

 Such small molecules mimetic agents may be produced by synthesizing a
25 plurality of peptides semi-peptidic compounds or non-peptidic, organic compounds, and then screening those compounds for their ability to inhibit the alpha4 integrin/alpha4 integrin ligand interaction. See generally U.S. Patent No. 4,833,092, Scott and Smith, "Searching for Peptide Ligands with an Epitope Library", Science, 249, pp. 386-90 (1990), and Devlin et al., "Random Peptide Libraries: A Source of Specific Protein
30 Binding Molecules", Science, 249, pp. 40407 (1990).

 In other preferred embodiments, the agent that is used in the method of the invention to bind to, including block or coat, cell-surface alpha4 integrin and/or alpha4 integrin ligand is an anti-VLA-4 and/or anti-alpha4beta7 monoclonal antibody or antibody homolog. Preferred antibodies and homologs for treatment, in particular for
35 human treatment, include human antibody homologs, humanized antibody homologs,

5 chimeric antibody homologs, Fab, Fab', F(ab')₂ and F(v) antibody fragments, and monomers or dimers of antibody heavy or light chains or mixtures thereof. Monoclonal antibodies against VLA-4 are a preferred binding agent in the method of the invention.

As used herein, the term "antibody homolog" includes intact antibodies consisting of immunoglobulin light and heavy chains linked via disulfide bonds. The term "antibody homolog" is also intended to encompass a protein comprising one or—
10 more polypeptides selected from immunoglobulin light chains, immunoglobulin heavy chains and antigen-binding fragments thereof which are capable of binding to one or more antigens. The component polypeptides of an antibody homolog composed of more than one polypeptide may optionally be disulfide-bound or otherwise covalently
15 crosslinked.

Accordingly, therefore, "antibody homologs" include intact immunoglobulins of types IgA, IgG, IgE, IgD, IgM (as well as subtypes thereof), wherein the light chains of the immunoglobulin may be of types kappa or lambda.

"Antibody homologs" also include portions of intact antibodies that retain antigen-binding specificity, for example, Fab fragments, Fab' fragments, F(ab')₂
20 fragments, F(v) fragments, heavy chain monomers or dimers, light chain monomers or dimers, dimers consisting of one heavy and one light chain, and the like. Thus, antigen-binding fragments, as well as full-length dimeric or trimeric polypeptides derived from the above-described antibodies are themselves useful.

As used herein, a "humanized antibody homolog" is an antibody homolog,
25 produced by recombinant DNA technology, in which some or all of the amino acids of a human immunoglobulin light or heavy chain that are not required for antigen binding have been substituted for the corresponding amino acids from a nonhuman mammalian immunoglobulin light or heavy chain.

As used herein, a "chimeric antibody homolog" is an antibody homolog,
30 produced by recombinant DNA technology, in which all or part of the hinge and constant regions of an immunoglobulin light chain, heavy chain, or both, have been substituted for the corresponding regions from another immunoglobulin light chain or heavy chain. In another aspect the invention features a variant of a chimeric molecule
35 which includes: (1) a VLA-4 targeting moiety, e.g., a VCAM-1 moiety capable of

5 binding to antigen (i.e., VLA-4) on the surface of VLA-4 bearing myeloma cells; (2) optionally, a second peptide, e.g., one which increases solubility or in vivo life time of the VLA-4 targeting moiety, e.g., a member of the immunoglobulin superfamily or fragment or portion thereof, e.g., a portion or a fragment of IgG, e.g., the human IgG1 heavy chain constant region, e.g., CH2 and CH3 hinge regions; and a toxin moiety. The VLA-4 targeting moiety can be any naturally occurring VLA-4 ligand or fragment
10 thereof, e.g., a VCAM-1 peptide or a similar conservatively substituted amino acid sequence. A preferred targeting moiety is a soluble VCAM-1 fragment, e.g., the N-terminal domains 1 and 2 of the VCAM-1 molecule. The chimeric molecule can be used to treat a subject, e.g., a human, at risk for disorder, e.g., multiple myeloma,
15 characterized by the presence of myeloma cells bearing VLA-4, and preferably activated VLA-4.

As used herein, a "human antibody homolog" is an antibody homolog produced by recombinant DNA technology, in which all of the amino acids of an immunoglobulin light or heavy chain that are derived from a human source.

20 Methods of Making Anti-VLA-4 Antibody Homologs

The technology for producing monoclonal antibody homologs is well known. Briefly, an immortal cell line (typically myeloma cells) is fused to lymphocytes (typically splenocytes) from a mammal immunized with whole cells expressing a given
25 antigen, e.g., VLA-4, and the culture supernatants of the resulting hybridoma cells are screened for antibodies against the antigen. See, generally, Kohler et al., 1975, *Nature*, 265: 295-297.

Immunization may be accomplished using standard procedures.. The unit dose and immunization regimen depend on the species of mammal immunized, its immune
30 status, the body weight of the mammal, etc. Typically, the immunized mammals are bled and the serum from each blood sample is assayed for particular antibodies using appropriate screening assays. For example, anti-VLA-4 antibodies may be identified by immunoprecipitation of 125I-labeled cell lysates from VLA-4-expressing cells. (See, Sanchez-Madrid et al. 1986, *Eur. J. Immunol.*, 16: 1343-1349 and Hemler et al. 1987, *J. Biol. Chem.*, 262, 11478-11485). Anti-VLA-4 antibodies may also be identified by
35

5 flow cytometry, e.g., by measuring fluorescent staining of Ramos cells incubated with an antibody believed to recognize VLA-4 (see, Elices et al., 1990 Cell, 60: 577-584). The lymphocytes used in the production of hybridoma cells typically are isolated from immunized mammals whose sera have already tested positive for the presence of anti-VLA-4 antibodies using such screening assays.

10 Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using 1500 molecular weight
15 polyethylene glycol ("PEG 1500"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridomas producing a desired antibody are detected by screening the hybridoma culture supernatants. For example, hybridomas prepared to produce anti-VLA-4
20 antibodies may be screened by testing the hybridoma culture supernatant for secreted antibodies having the ability to bind to a recombinant alpha4-subunit-expressing cell line (see, Elices et al., supra).

To produce anti-VLA-4 antibody homologs that are intact immunoglobulins, hybridoma cells that tested positive in such screening assays were cultured in a nutrient
25 medium under conditions and for a time sufficient to allow the hybridoma cells to secrete the monoclonal antibodies into the culture medium. Tissue culture techniques and culture media suitable for hybridoma cells are well known. The conditioned hybridoma culture supernatant may be collected and the anti-VLA4 antibodies optionally further purified by well-known methods.

30 Alternatively, the desired antibody may be produced by injecting the hybridoma cells into the peritoneal cavity of an unimmunized mouse. The hybridoma cells proliferate in the peritoneal cavity, secreting the antibody which accumulates as ascites fluid. The antibody may be harvested by withdrawing the ascites fluid from the peritoneal cavity with a syringe.

5 Several mouse anti-VLA-4 monoclonal antibodies have been previously described. See, e.g., Sanchez-Madrid et al., 1986, supra; Hemler et al., 1987, supra; Pulido et al., 1991, J. Biol. Chem., 266 (16), 10241-10245). These anti-VLA-4 monoclonal antibodies such as HP 1/2 and other anti-VLA-4 antibodies (e.g., HP2/1, HP2/4, L25, P4C2, P4G9) capable of recognizing the P chain of VLA-4 will be useful
10 in the methods of treatment according to the present invention. AntiVLA-4 antibodies that will recognize the VLA-4 alpha4 chain epitopes involved in binding to VCAM-1 and fibronectin ligands (i.e., antibodies which can bind to VLA-4 at a site involved in ligand recognition and block VCAM-1 and fibronectin binding) are preferred. Such antibodies have been defined as B epitope-specific antibodies (B1 or B2) (Pulido et al.,
15 1991, supra) and are also anti-VLA-4 antibodies according to the present invention.

 Fully human monoclonal antibody homologs against VLA-4 are another preferred binding agent which may block or coat VLA-4 antigens in the method of the invention. In their intact form these may be prepared using in vitro-primed human splenocytes, as described by Boerner et al., 1991, J. Immunol., 147, 86-95.
20 Alternatively, they may be prepared by repertoire cloning as described by Persson et al., 1991, Proc. Nat. Acad. Sci. USA, 88: 2432-2436 or by Huang and Stollar, 1991, J. Immunol. Methods 141, 227-236. U.S. Patent 5,798,230 (Aug. 25, 1998, "Process for the preparation of human monoclonal antibodies and their use") who describe preparation of human monoclonal antibodies from human B cells. According to this
25 process, human antibody-producing B cells are immortalized by infection with an Epstein-Barr virus, or a derivative thereof, that expresses Epstein-Barr virus nuclear antigen 2 (EBNA2). EBNA2 function, which is required for immortalization, is subsequently shut off, which results in an increase in antibody production.

 In yet another method for producing fully human antibodies, United States
30 Patent 5,789,650 (Aug. 4, 1998, "Transgenic non-human animals for producing heterologous antibodies") describes transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. Endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against
35 endogenous immunoglobulins. Heterologous antibodies are encoded by

5 immunoglobulin genes not normally found in the genome of that species of non-human animal. One or more transgenes containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of functionally rearranging transgenic immunoglobulin sequences and producing a repertoire of antibodies of various isotypes
10 encoded by human immunoglobulin genes. Such heterologous human antibodies are-- produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous, fully human antibody homolog.

15 Large nonimmunized human phage display libraries may also be used to isolate high affinity antibodies that can be developed as human therapeutics using standard phage technology (Vaughan et al, 1996). Yet another preferred binding agent which may block or coat VLA-4 antigens in the method of the invention is a humanized recombinant antibody homolog having anti-VLA-4 specificity. Following the early
20 methods for the preparation of chimeric antibodies, a new approach was described in EP 0239400 (Winter et al.) whereby antibodies are altered by substitution of their complementarity determining regions (CDRs) for one species with those from another. This process may be used, for example, to substitute the CDRs from human heavy and light chain Ig variable region domains with alternative CDRs from murine variable
25 region domains. These altered Ig variable regions may subsequently be combined with human Ig constant regions to created antibodies which are totally human in composition except for the substituted murine CDRs. Such CDR-substituted antibodies would be predicted to be less likely to elicit an immune response in humans compared to chimeric antibodies because the CDR-substituted antibodies contain considerably
30 less non-human components. The process for humanizing monoclonal antibodies via CDR "grafting" has been termed "reshaping". (Riechmann et al., 1988, Nature 332, 323-327; Verhoeven et al., 1988, Science 239, 1534-1536).

Typically, complementarity determining regions (CDRs) of a murine antibody are transplanted onto the corresponding regions in a human antibody, since it is the
35 CDRs (three in antibody heavy chains, three in light chains) that are the regions of the

5 mouse antibody which bind to a specific antigen. Transplantation of CDRs is achieved by genetic engineering whereby CDR DNA sequences are determined by cloning of murine heavy and light chain variable (V) region gene segments, and are then transferred to corresponding human V regions by site directed mutagenesis. In the final stage of the process, human constant region gene segments of the desired isotype
10 (usually gamma I for CH and kappa for CL) are added and the humanized heavy and – light chain genes are co-expressed in mammalian cells to produce soluble humanized antibody.

The transfer of these CDRs to a human antibody confers on this antibody the antigen binding properties of the original murine antibody. The six CDRs in the murine
15 antibody are mounted structurally on a V region “framework” region. The reason that CDR-grafting is successful is that framework regions between mouse and human antibodies may have very similar 3-D structures with similar points of attachment for CDRs, such that CDRs can be interchanged. Such humanized antibody homologs may be prepared, as exemplified in Jones et al., 1986, Nature 321, 522-525; Riechmann,
20 1988, Nature 332, 323-327; Queen et al., 1989, Proc. Nat. Acad. Sci. USA 86, 10029; and Orlandi et al., 1989, Proc. Nat. Acad. Sci. USA 86, 3833.

Nonetheless, certain amino acids within framework regions are thought to interact with CDRs and to influence overall antigen binding affinity. The direct transfer of CDRs from a murine antibody to produce a recombinant humanized antibody without
25 any modifications of the human V region frameworks often results in a partial or complete loss of binding affinity. In a number of cases, it appears to be critical to alter residues in the framework regions of the acceptor antibody in order to obtain binding activity

Queen et al., 1989 (supra) and WO 90/07861 (Protein Design Labs) have
30 described the preparation of a humanized antibody that contains modified residues in the framework regions of the acceptor antibody by combining the CDRs of a murine MAb (anti-Tac) with human immunoglobulin framework and constant regions. They have demonstrated one solution to the problem of the loss of binding affinity that often results from direct CDR transfer without any modifications of the human V region
35 framework residues; their solution involves two key steps. First, the human V

5 framework regions are chosen by computer analysts for optimal protein sequence
homology to the V region framework of the original murine antibody; in this case, the
anti-Tac MAb. In the second step, the tertiary structure of the murine V region is
modelled by computer in order to visualize framework amino acid residues which are
likely to interact with the murine CDRs and these murine amino acid residues are then
10 superimposed on the homologous human framework. See also Protein Design Labs –
U.S. Patent 5,693,762.

One may use a different approach (Tempest et al., 1991, Biotechnology 9, 266-
271) and utilize, as standard, the V region frameworks derived from NEWM and REI
heavy and light chains respectively for CDR-grafting without radical introduction of
15 mouse residues. An advantage of using the Tempest et al., approach to construct
NEWM and REI based humanized antibodies is that the 3dimensional structures of
NEWM and REI variable regions are known from x-ray crystallography and thus
specific interactions between CDRs and V region framework residues can be modeled.

Regardless of the approach taken, the examples of the initial humanized
20 antibody homologs prepared to date have shown that it is not a straightforward process.
However, even acknowledging that such framework changes may be necessary, it is not
possible to predict, on the basis of the available prior art, which, if any, framework
residues will need to be altered to obtain functional humanized recombinant antibodies
of the desired specificity. Results thus far indicate that changes necessary to preserve
25 specificity and/or affinity are for the most part unique to a given antibody and cannot be
predicted based on the humanization of a different antibody.

Preferred antagonists useful in the present invention include chimeric
recombinant and humanized recombinant antibody homologs (i.e., intact
immunoglobulins and portions thereof) with B epitope specificity that have been
30 prepared and are described in co-pending U.S. Patent Application Serial No.
08/004,798, filed January 12, 1993, PCT Publication US94/00266, filed January 7,
1994. The starting material for the preparation of chimeric (mouse V - human C) and
humanized anti-VLA-4 antibody homologs may be a murine monoclonal anti-VLA-4
antibody as previously described, a monoclonal anti-VLA-4 antibody commercially
35 available (e.g., HP2/1, Amac International, Inc., Westbrook, Maine), or a monoclonal

5 anti-VLA-4 antibody prepared in accordance with the teaching herein. For example, the variable regions of the heavy and light chains of the anti-VLA-4 antibody HP ½ have been cloned, sequenced and expressed in combination with constant regions of human immunoglobulin heavy and light chains. Such HP ½ antibody is similar in specificity and potency to the murine HP 1/2 antibody, and may be useful in methods
10 of treatment according to the present invention. —

Other preferred humanized anti-VLA4 antibody homologs are described by Athena Neurosciences, Inc. in PCT/US95/01219 (27 July 1995) These humanized anti-VLA-4 antibodies comprise a humanized light chain and a humanized heavy chain. The humanized light chain comprises three complementarity determining regions (CDR1,
15 CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of a mouse 21-6 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence except in at least position the amino acid position is occupied by the same amino acid present in the equivalent position of the mouse 21.6 immunoglobulin light
20 chain variable region framework. The humanized heavy chain comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of a mouse 21-6 immunoglobulin heavy chain, and a variable region framework from a human heavy chain variable region framework sequence except in at least one position the amino acid
25 position is occupied by the same amino acid present in the equivalent position of the mouse 21-6 immunoglobulin heavy chain variable region framework.

Therapeutic Applications

In this method according to the first aspect of the invention, VLA-4 binding
30 agents, in particular, VCAM fusions and anti-VLA-4 antibody homologs are preferably administered parenterally. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques

The VLA-4 binding agents are preferably administered as a sterile
35 pharmaceutical composition containing a pharmaceutically acceptable carrier, which

5 may be any of the numerous well known carriers, such as water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, and the like, or combinations thereof. The compounds of the present invention may be used in the form of pharmaceutically acceptable salts derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, tris(hydroxymethyl)methylamine and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

The pharmaceutical compositions of this invention comprise any of the compounds of the present invention, or pharmaceutically acceptable derivatives thereof, together with any pharmaceutically acceptable carrier. The term "carrier" as used herein includes acceptable adjuvants and vehicles. Pharmaceutically acceptable carriers that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium

5 trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

According to this invention, the pharmaceutical compositions may be in the form of a sterile injectable preparation, for example a sterile injectable aqueous or
10 oleaginous suspension. This suspension may be formulated according to techniques — known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are
15 water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as do natural pharmaceutically-acceptable oils, such as olive oil or castor
20 oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as Ph. Helv or similar alcohol.

The pharmaceutical compositions of this invention, in particular small molecule antagonists of the VLA-4/VCAM-1 interaction, may be given parenterally or orally. If
25 given orally, they can be administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch.
30 When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added. Topically-transdermal patches may also be used. The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation through the use of a nebulizer, a dry powder inhaler or a metered
35 dose inhaler. Such compositions are prepared according to techniques well-known in

5 the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

10 According to another embodiment compositions containing a compound of this invention may also comprise an additional agent selected from the group consisting of corticosteroids, antiinflammatories, immunosuppressants, antimetabolites, and immunomodulators. Specific compounds within each of these classes may be selected from any of those listed under the appropriate group headings in "Comprehensive Medicinal Chemistry", Pergamon Press, Oxford, England, pp. 970-986 (1990), the disclosure of which is herein incorporated by reference. Also included within this group are compounds such as theophylline, sulfasalazine and aminosalicylates (antiinflammatories); cyclosporin, FK-506, and rapamycin (immunosuppressants); cyclophosphamide and methotrexate (antimetabolites); steroids (inhaled, oral or topical) and interferons (immunomodulators).

20 The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, and the particular mode of administration. It should be understood, however, that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of active ingredient may also depend upon the therapeutic or prophylactic agent, if any, with which the ingredient is co-administered.

30 The dosage and dose rate of the compounds of this invention effective to prevent, suppress or inhibit cell adhesion will depend on a variety of factors, such as the nature of the inhibitor, the size of the patient, the goal of the treatment, the nature of the pathology to be treated, the specific pharmaceutical composition used, and the judgment of the treating physician. Dosage levels of between about 0.001 and about 100 mg/kg body weight per day, preferably between about 0.1 and about 50 mg/kg body weight per day of the active ingredient compound are useful. Most preferably, the VLA-4 binding

35

applications, each consisting of generally the same disclosure, and each being subjected to essentially the same search, perhaps by different Examiners on different occasions. This process would place an unnecessary burden on both the Patent and Trademark Office and on the Applicants.

Regardless of whether the five inventions are independent or distinct, Applicants respectfully assert that the Examiner need not have restricted the application. MPEP § 803 requires that "[i]f the search and examination of an entire application can be made without serious burden, the Examiner must examine it on the merits, even though it includes claims to independent or distinct inventions." Therefore, it is not mandatory to make a restriction requirement in all situations where it would be deemed proper.

In the interest of economy, for the Office, for the public-at-large, and for Applicants, reconsideration and withdrawal of the restriction requirement are requested.

Nevertheless, in order to comply with the requirements of 37 C.F.R. § 1.143, Applicants indicate below an election of one group for examination. As such, Applicants elect, with traverse, to prosecute the invention of Group I, namely claims 1-24 and 46-59, drawn to a method of promoting remyelination of nerve cells or reversing paralysis in a mammal comprising administering a remyelinating agent, for prosecution in the above-identified application.

The Examiner also requires identification of a single disclosed species under 35 U.S.C. § 121. As Applicants have elected Group I, the Office requires the election of a species. With regard to this election of species, Applicants elect, with traverse, natalizumab or immunologically active fragment thereof as the agent, and multiple sclerosis as the condition, for the purposes of searching only. It is believed that claims 1-24 and 46-59 within the elected group (Group I) are readable upon the elected species of natalizumab or immunologically active fragment thereof as defined above. It is believed that claims 1-4 and 46-47 within the elected group (Group I) are readable upon the elected species of multiple sclerosis thereof as defined above.

5 agent, if an antibody or antibody derivative, will be administered at a dose ranging
between about 0.1 mg/kg body weight/day and about 20 mg/kg body weight/day,
preferably ranging between about 0.1 mg/kg body weight/day and about 10 mg/kg body
weight/day and at intervals of every 1-14 days. For non-antibody or small molecule
binding agents, the dose range should preferably be between molar equivalent amounts
10 to these amounts of antibody. Preferably, an antibody composition is administered in
an amount effective to provide a plasma level of antibody of at least 1 mg/ml.
Optimization of dosages can be determined by administration of the binding agents,
followed by assessment of the coating of VLA-4-positive cells by the agent over time
after administered at a given dose in vivo.

15 Myeloma cells contained in a sample of the individual's peripheral blood (or
bone marrow cells) should be probed for the presence of the agent in vitro (or ex vivo)
using a second reagent to detect the administered agent. For example, this may be a
fluorochrome labelled antibody specific for the administered agent which is then
measured by standard FACS (fluorescence activated cell sorter) analysis. Alternatively,
20 presence of the administered agent may be detected in vitro (or ex vivo) by the inability
or decreased ability of the individual's cells to bind the same agent which has been
itself labelled (e.g., by a fluorochrome). The preferred dosage should produce
detectable coating of the vast majority of VLA-4-positive cells. Preferably, coating is
sustained in the case of an antibody homolog for a 1- 14 day period.

25 Animal Models:

The animal model has been described in detail (Garrett 1997). Briefly, Radl et al
(1988) had described a murine model of myeloma which arose spontaneously in aged
C57BL/KaLwRij mice. This condition occurred in approximately 1 in 200 animals as
30 they aged, and led to a monoclonal gammopathy with some of the features of human
disease (Radl 1988). To develop a better and more reproducible animal model we have
established and subcloned a cell line from this murine myeloma called 5TGM1, and
found that it causes lesions in mice characteristic of human myeloma, such as severe
osteolysis and the involvement of non-bone organs including liver and kidney (Garrett
35 1997). Mice inoculated with cultured cells develop disease in a highly predictable and

5 reproducible manner, which includes formation of a monoclonal gammopathy and
radiologic bone lesions. Furthermore, some of the mice become hypercalcemic, and the
bone lesions are characterized by increased osteoclast activity. Thus, based on
histological examination of affected organs in 5TGM1-bearing mice and increased
serum levels of IgG2b, 5TGM1 is defined as a murine myeloma which recapitulates
10 accurately the hallmarks of human disease. —

 The following examples are intended to further illustrate certain preferred
embodiments of the invention and are not intended to be limiting in nature. In the
following examples, the necessary restriction enzymes, plasmids, and other reagents
15 and materials may be obtained from commercial sources and cloning, ligation and other
recombinant DNA methodology may be performed by procedures well-known in the
art.

20 Example 1: MATERIALS AND METHODS

5TGM1 Myeloma Cells

 5TGM1 myeloma cells were initially derived from a myeloma which arose
spontaneously in aged C57BL/KaLwRij mice (Garrett 1997, Vanderkerken 1997).
25 Cells were grown in Isocove's Modified Dulbecco's Medium (IMDM, Life
Technologies Inc., Gaithersburg, MD) supplemented with 10% fetal bovine serum
(FBS, Summit, Fort Collins, CO) and 1 % penicillin-streptomycin solution (GIBCO,
Grand Island, NY) at 37 C in 5% CO₂ atmosphere. For in vitro experimentation
described below, 5TGM1 cells between passage 25 and 30 were used.

30

Antibodies, soluble VCAM-1

 Neutralizing antibodies against murine VCAM-1 (M/K-2.7), integrin VLA-4
(PS/2), and Intercellular Adhesion Molecule-1 (ICAM-1, YN1/1.7), were kindly gifted
by Dr. Kensuke Miyake (Saga Medical University, Saga, Japan). Recombinant soluble

5 VCAM-1 (Lobb et al, 1991), containing the 7 extracellular domains of human VCAM-1, was the gift of Dr. Roy Lobb, Biogen Inc., Cambridge, MA.

10 Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

Using RT-PCR, we confirmed expression of VCAM-1 and integrin alpha4 in bone marrow stromal cells and 5TGM1, respectively. Total RNA was prepared from 5TGM1, a primary culture of bone marrow stromal cells and an ST2 marrow stromal cell line (RIKEN Cell Bank, Tsukuba, Japan) by the single-step RNA isolation method using TRIzol reagent (GIBCO). Three μ g of RNA was incubated with 50 ng of random hexamer at 70 ° C for 10 min and chilled on ice, then converted to first strand cDNA using reverse transcriptase (Perkin-Elmer, Branchburg, NJ) according to the manufacturers instruction. The primers used for PCR were as follows: murine VCAM-1 5'-primer; 5'-OH-GCTGCGCGTCACCATTGTTCTC-3'-OH [SEQ ID NO: 1];
20 murine VCAM-1 3'-primer; 5'-OH-ACCACCCTCTTGAAGCCTTGTG-3'-OH [SEQ ID NO: 2] ; murine integrin alpha4 5'- primer; 5'-OH-CCCCCTCAACACGAACAGATAGG-3'-OH [SEQ ID NO: 3]; murine integrin alpha4 3'-primer; 5'-OH-GCCTTGTCCTTAGCAACACTGC-3'-OH [SEQ IDNO: 4].

PCR was performed for 30 cycles consisting of 1 min at 94° C, 1 min at 55° C
25 and 2 min at 72° C. PCR reaction mixture (total 50 μ l) contained 10 microliters. First strand cDNA, 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 2 mM MgCl₂, deoxy-NTP mix (0.2 mM each), the pair of primers (0.15 micromolar each) and 2 U Taq DNA polymerase (Perkin-Elmer, Branchburg, NJ). The PCR products were separated on 2.5% agarose gels containing ethidium bromide and visualized under ultraviolet light.
30 The size of the fragments were confirmed by reference to molecular weight markers.

Attachment of 5TGM1 Cells onto Bone Marrow Stromal Cells

For heterotypic cell-cell adhesion assays, ST2 cells (5 e 4/well) were seeded in 48-well culture plates (Costar, Cambridge, MA) and cultured 48 h in alphaMEM
35 supplemented with 10% FBS until confluency. 5TGM1 cells (5 e 6) were labeled by

5 incubation with 10 microCi [methyl- ^3H] thymidine (New England Nuclear) for 24 h at
37° C in the culture medium. After the ST2 monolayer was formed, it was incubated
with 1 % bovine serum albumin (BSA, Sigma, St Louis, MO) in serum-free alphaMEM
for 1 hours and tritium-labeled 5TGM1 cells were plated onto the monolayer. The
system was incubated in the absence or presence of antibodies to VCAM-1 or
10 alpha4beta1 integrin at 37° C for 1 h. Non-adherent cells were removed by washing –
with 5% trichloroacetic acid twice and PBS twice, and adherent cells were solubilized
in 300 microliters of 0.25 mM NaOH, neutralized with the same volume of 0.25 mM
HCl and the radioactivity was determined in a liquid scintillation counter.

15 **Osteoclast Formation Assay in the Co-culture of 5TGM1 and Mouse Bone Marrow Cells**

Mouse bone marrow cells were obtained from 5-week-old male C57BL mice as
described previously (Yoneda 1993). Femurs and tibiae were dissected aseptically and
both ends cut off. Bone marrow cells were flushed out, collected and incubated in
20 alphaMEM supplemented with 10% FBS (Hyclone, Logan, UT) and 1 % penicillin-
streptomycin in 100 mm-culture dishes (Becton Dickinson Labware, Bedford, MA) at
37° C for 2 h. Non-adherent cells containing hemopoietic osteoclast precursors and
stromal cells were harvested. Bone marrow cells (1 e 6) and 5TGM1 cells (1 e 3) in
300 microliters of the culture medium were plated onto 48-well culture plates (day 0).
25 On day 2, 300 microliters of fresh culture medium was gently added to each well, and
on day 4, 300 microliters of spent medium was replaced with the same volume of fresh
medium. On day 6, the cultures were fixed and stained for tartrate-resistant acid
phosphatase (TRAP) using commercial kits (Sigma). TRAP-positive multinucleated
cells with more than 3 nuclei were defined as osteoclast-like (OC-like) cells, and
30 manually counted under microscope. To confirm that these OC-like cells have the
capability to resorb bone, 5TGM1 cells and marrow cells were co-cultured on 5x5 mm
whale dentine slices in the same condition, and resorption pits formed on these dentine
slices were examined by scanning electron microscopy as described (Yoneda 1992).

In some experiments, co-cultures of 5TGM1 myeloma cells and marrow cells
35 were performed using transwell inserts (Becton Dickinson Labware) to prevent direct

5 contact between these two types of cells. (2 e 6 , 24-well plates, Costar). Marrow cells were plated in the lower chambers and 5TGM 1 myeloma cells (2e 3) were then plated in either lower (direct contact) or upper (no contact) chambers.

Organ Cultures of ⁴⁵Ca-labelled Fetal Rat Long Bones

10 Conditioned media harvested from 5TGM1 cultures were assayed for bone- – resorbing activity by organ cultures of ⁴⁵Ca-labelled fetal rat long bones as described previously (Mbalaviele 1995). Pregnant rats were injected with 250 uCi of ⁴⁵Ca (New England Nuclear) on the 18th day of gestation. Radius and ulna bone shafts were obtained from 19-day fetuses by microdissection, and precultured for 24 h in BGJ
15 medium (Sigma) supplemented with 0.1 % BSA between air and liquid-phase on stainless mesh grids. Bones were then cultured in the presence of conditioned media (50% v/v) or in control medium for 120 hours. The media were changed once at 48 hours. At the end of the culture, bones were incubated in ice-cold 5% trichloroacetic acid for 2h , and ⁴⁵Ca radioactivity in bones and media determined in a liquid
20 scintillation counter. Bone resorption was quantitated as the percentage of ⁴⁵Ca released into the medium from bones as calculated by: (⁴⁵Ca count in medium)/ (⁴⁵Ca count in medium and bone) x 100.

Co-culture of 5TGM1 Myeloma Cells with Mouse Stromal Cell Line ST2 Cells

25 ST2 cells (0.5 e 6) and 5TGM1 (4 e 6) cells were plated together onto 60-mm culture dishes (Beckton Dickinson) in 10% FBS-supplemented IMDM and cultured overnight, washed with serum-free IMDM twice, and incubated in 5 ml of serum-free IMDM. After 48 h, conditioned media were harvested and stored at -70 ° C until use.

Effect of mAb PS2 to VLA-4 on serum IgG2b elevation in 5TGM1-bearing mice

30 Mice were injected with 1 e 5 5TGM1 cells, which were allowed to colonize the bone marrow. Mice were split into two groups of three, one serving as a control group, and the second treated biweekly beginning on day 8 with 80 ug mAb PS/2 (4 mg/kg). Levels

5 of IgG2b, the antibody isotype produced by 5TGM1 myeloma cells, were measured weekly from weeks 1 to 6.

RESULTS

10

Expression of VCAM-1, VLA-4, and effect of Antibodies Against VCAM-1 and VLA-4 on 5TGM1 Attachment to ST2 Monolayers

15

20

Using RT-PCR, we confirmed the expression of VCAM-1 and integrin VLA-4 in bone marrow stromal cells and myeloma cells, respectively. As expected, both the ST2 stromal cell line and primary bone marrow stromal cells expressed VCAM-1, while 5TGM1 did not. In contrast, the 5TGM1 myeloma cells expressed integrin VLA-4, whereas stromal cells did not (data not shown). In addition, both anti-VCAM-1 antibody (10 ug/ml) and VLA-4 antibody (10 ug/ml) partially (50-80%) inhibited the attachment of 5TGM1 cells to ST2 monolayers, showing that VCAM-1 and the VLA-4 integrin expressed on these cells are biologically functional and that these antibodies have neutralizing activity (data not shown).

25

OC-like Cell Formation in the Coculture of 5TGM1 Myeloma Cells with Mouse Bone Marrow Cells

30

On day 6 of the coculture of 5TGM1 cells and mouse marrow cells, numerous TRAP-positive multinucleated osteoclast-like (OC-like) cells were formed. These OC-like cells exhibited resorption pit formation on dentine slices, demonstrating that these cells were capable of resorbing bone, and possess an osteoclastic phenotype. In experiments using transwell inserts, formation of OC-like cells was observed when 5TGM1 cells were cultured in direct contact with bone marrow cells. In contrast, there was only a marginal number of OC-like cells formed when 5TGM1 cells were separated from marrow cells by the transwell membrane. Thus 5TGM1 cells induce osteoclast formation in mixed marrow cultures, and this induction requires direct cell-cell contact.

35

Effect of Antibodies Against VCAM-1 and Integrin VLA4 on OC-like Cell Formation in the Co-culture of 5TGM1 and Marrow Cells

5 Both anti-VCAM-1 antibody (VCAM-1 Ab, 10ug/ml) and anti VLA-4-integrin antibody (alpha4beta1 Ab, 10 ug/ml) dramatically inhibited OC-like cell formation. In contrast mAb against ICAM-1, another adhesion molecule on marrow stromal cells implicated in stromal/myeloma interactions, had no effect on OC-like cell formation (Figure 1).

10 To determine whether this inhibition by VCAM-1 and VLA-4 mAbs was specific for 5TGM1-induced OC-like cell formation and was not due to cytotoxicity, the effects of these antibodies were examined on OC-like cell formation induced by 1,25 (OH)₂ D₃, a widely-used stimulator of osteoclastogenesis in mouse bone marrow cell cultures (Takahashi 1988). Neither VCAM-1 Ab nor VLA-4 mAb inhibited OC-like
15 cell formation induced by vitamin D₃, which itself had no effect on VCAM-1 expression in stromal cells (data not shown).

Effect of conditioned Medium Harvested from the Co-culture of 5TGM1 and ST2 on Bone Resorption

20 Conditioned medium from the co-culture of 5TGM1 cells and ST2 cells showed a marked increase in bone resorption in the fetal rat long bone assay (Figure 2), while conditioned medium of 5TGM1 caused only a marginal increase, s compared to control medium. Conditioned medium from ST2 cells showed no increase in bone resorption. Thus direct cell-cell contact via VCAM-1 and VLA-4 both induces osteoclast-like cells
25 and production of bone-resporboing factors in vitro.

Effect of Recombinant Soluble VCAM-1 (sVCAM-1) on the Production of Bone-resorbing and Osteoclastogenic Activity by 5TGM1 Cells

30 Conditioned medium of 5TGM1 treated with a soluble recombinant form of VCAM-1 (sVCAM-1) increased bone resorption in fetal rat long bones in a dose-dependent manner, while conditioned medium obtained from untreated 5TGM1 only marginally increased bone resorption. Soluble VCAM-1 itself had no effects on bone resorption (data not shown). In the mouse marrow culture system, conditioned medium harvested from 5TGM1 cells treated with sVCAM-1 showed increased activity of OC-

5 like cell formation, while conditioned medium of untreated 5TGM1 exhibited only marginal activity of OC-like cell formation (Figure 3).

Expression of Rank ligand mRNA in marrow stromal cells (ST2) cultured in the presence and absence of murine myeloma cells

10

Because Rank ligand appears to be an important mediator of OCL formation and may be the final common pathway for the effects of osteoclastogenic cytokines on OCL formation, we have examined the expression of Rank ligand in 5TGM1 and ST2 cells both individually and when cocultured. We find that coculture of 5TGM1 and ST2 cells induces Rank ligand mRNA in the ST2 cells. Furthermore, while 5TGM1 cells do not express Rank ligand, they do so when treated with sVCAM-1 (not shown). Finally, the conditioned medium from 5TGM1 cells treated with sVCAM-1 induced Rank ligand mRNA in ST2 cells, suggesting that the VCAM-1/VLA-4 pathway produces a cytokine in myeloma cells that enhances Rank ligand expression by marrow stromal cells (data not shown).

20

In summary, we show that 5TGM1 cells alone produce marginal amount of activity that stimulates OC-like cell formation and bone resorption. However, when 5TGM1 myeloma cells were co-cultured with bone marrow cells containing hemopoietic osteoclast precursors and stromal cells, they strongly adhered to the stromal cells and increased OC-like cell formation. There were no OC-like cells formed in the co-cultures in which 5TGM1 cells were prevented from contacting stromal cells. Furthermore, in organ cultures of fetal rat long bones the conditioned medium harvested from the co-cultures of 5TGM1 myeloma cells and ST2 bone marrow stromal cells had increased bone resorbing activity compared with conditioned medium of either ST2 or 5TGM1 alone. These data are consistent with the notion that direct cell-cell contact of 5TGM1 cells with bone marrow stromal cells is required for the production of osteoclast-stimulating and bone-resorbing activity. We then determined what cell adhesion molecules were involved in the direct cell-cell interaction between 5TGM1 cells and marrow stromal cells that is necessary for the production of osteoclastogenic activity.

30

35

5 Our data indicate that VCAM-1 and VLA-4 integrin play a role in this cell-cell interaction, since neutralizing antibodies to these two adhesion molecules profoundly decreased OC-like cell formation in the co-cultures. The VCAM-1/VLA-4 integrin interaction is responsible for the cell-cell communication between marrow stromal cells and 5TGM1 myeloma cells leading to increased production of a osteoclastogenic and
10 bone-resorbing activity. Finally, this bone resorbing activity in part is due to induction of Rank ligand.

Example 2: *IN VIVO* EXPERIMENTS

15 Our in vitro studies suggest that the interaction between VLA-4 on myeloma cells with VCAM-1 on marrow stromal cells may play a key role in the induction of bone resorbing activity by myeloma. We have taken the key step of testing this hypothesis in vivo in an animal model which accurately reflects human disease.

20 A. In this experiment, mice were injected with 1×10^5 5TGM1 myeloma cells, which were allowed to colonize the bone marrow. Mice were split into two groups of three, one serving as a control group, and the second treated biweekly beginning on day 8 with mAb PS/2. Levels of IgG2b, the antibody isotype produced by 5TGM1 myeloma cells, were measured weekly from weeks 1 to 6. Treatment with mAb at a dose of 80 ug
25 per injection (~ 4 mg/kg) biweekly strongly inhibited IgG2b production, indicative of significant inhibition of myeloma cell survival and growth in vivo (Figure 4). Further, the treated mice showed reduced incidence of paraplegia (all 3 untreated animals showed paraplegia on day 42, while only one of the treated animals showed paraplegia. The two treated animals with no paraplegia also showed a reduction in spleen and liver
30 weights, which also correlate with tumor burden. Finally, the treated animals showed a reduction in tumor area by histology (from 6.71 ± 1.74 to 0.05 ± 0.08 square millimeters) in the tibia and femurs. There was no effect of treatment on serum calcium levels (data not shown)

5 B. In a parallel experiment, treatment with 40 ug PS/2 biweekly had no effect on IgG2b levels (not shown). These data show that mAb PS/2 to VLA-4 strongly inhibits the growth of established myeloma cells in a dose-dependent fashion.

10 C. In another in vivo experiment, 18 SCID mice were injected with 5TGM1 myeloma cells at day 0. Four mice were treated with PBS; 4 mice were treated in a prophylactic protocol with mAb M/K-2.7 reactive against to mouse VCAM1 at a dosage of 80 ug (- 4 mg/kg) every 3 days starting at day -1 (i.e. days - 1, 2, 5, 8, and 11). In a parallel experiment using the same protocol, five mice were treated with 160 ug mAb M/K-2.7 . In addition, five mice were treated with 160 ug mAb M/K-2.7
15 starting at day 8 (i.e. days 8, 11, 14, 17, and 20) in a therapeutic protocol. Serum was taken from all mice on days 21, 28, and 35, and animals were X-rayed then sacrificed for histology on day 35. All three treatment groups showed a reduction in serum IgG2b levels, indicative of reduced myeloma cell burden (Figure 5). A significant effect was also observed on spleen weights at the low dose prophylactic protocol relative to control
20 (0.23 +/- 0.14 g for control versus 0.08 +/- 0.04 for treated). In the prophylactic high dose group 4 of 5 animals showed a clear reduction in spleen weight, but the overall value was not significant because of one animal with a large spleen weight (data not presented).

25 D. One can investigate whether an initial high bolus dose of alpha4 integrin antagonist, followed by a maintenance dose, improves efficacy. The myeloma cells are already established in the marrow compartment, and their tight VLA-4-dependent interaction with VCAM-1 needs to be inhibited. Furthermore, presumably the greater the number of established myeloma cells, the higher the initial dose required to flush
30 cells out into the peripheral circulation.

 A larger study with the anti-VLA-4 antibody PS/2 was therefore performed. Twenty eight SCID mice were injected with 5TGM1 myeloma cells at day 0. Nine mice received no treatment; 9 mice received an isotype-matched control IgG mAb; 10 mice were treated with mAb PS/2 to alpha 4 integrin. A different therapeutic regimen

5 was given, in which mice were given a high dose of mAb (200 ug) on days 4,5, and 6, then a maintenance dose of 80 ug (- 4 mg/kg) every 3 days starting at day 8.

There was a statistically significant reduction in serum IgG2b when the treated group was compared to either the untreated or control IgG-treated group at weeks 3 and 4 (data not presented). Importantly, when the treated group was compared to either the untreated or control
10 IgG-treated group there was a clear effect on survival (Figure 6).

Example 3: OTHER IN VIVO EXPERIMENTS

Based on the information presented herein for the first time, persons having ordinary skill in the art can readily confirm and extend the importance of the alpha4
15 integrins and their ligands in multiple myeloma using the murine animal model described.

The following series of experiments are well within the level of skill in the art based upon the present disclosure but serve merely to exemplify, and not limit, the types of work.

20

- 1) Dose response to mAb PS/2 to determine the optimal biweekly maintenance dose. 80 ug shows good efficacy, but 40 ug was without effect. One examines higher doses up to 20 mg/kg two or three times weekly to determine optimal dosing.
- 2) Patients present with disease at different stages of severity, linked to increased
25 tumor burden. One examines the efficacy of mAb PS/2 given at different times after establishment of disease, i.e. one compares treatment initiation at 8 days (see for example Figure 4) to initiation after two, three, four and five weeks post inoculation to see how late mAb can be given to provide some relief of symptoms.
- 3) The effects of mAb MK-2 to murine VCAM-1 are examined, following the same
30 parameters outlined above (dosing, timing of dosing) for mAb to VLA-4. It is anticipated that similar dosing levels will be required to see efficacy.
- 4) Further markers of myeloma progression are examined, including tumor burden in both marrow and extramedullary sites, quantification of bone lesions by radiographic analysis of the skeleton by histomorphometry; measurement of rates of bone
35 reportation by evaluation of collagen crosslinks in plasma; measurement of

- 5 monoclonal protein production in plasma; hypercalcemia where present; and mortality.
- 5) Multiple myeloma is currently treated inefficiently with standard chemotherapeutic regimens. The additive or synergistic effects of mAbs at optimal dosing in conjunction with, or either before or after, dosing with appropriate
- 10 chemotherapeutic regimens is examined.
- 6) The ability of a small molecule alpha4 integrin inhibitor that is selective for one particular alpha4 integrin or is selective for several alpha4 integrins at once or the ability of combinations of such inhibitors, to mimic the effects of mAbs and block myeloma progression is examined using the protocols and outcomes described
- 15 above. Small molecule inhibitors are delivered parenterally or orally, in the dosing range of 0.1 to 30 mg/kg, once or twice daily, or twice or three times weekly.

Additional References:

- 20 Alsina M, Boyce B, Devlin R, Anderson JL, Craig F, Mundy GR, Roodman GD. Development of an in vivo model of human multiple myeloma bone disease. *Blood* 87: 1495-1501, 1996.
- 25 Attal M, Harousseau JL, Stoppa AM, Sotto JJ, Fuzibet JG, Rossi JF, Casassus P, Maisonneuve H, Facon T, Ifrah N, Payen C, Bataille R. A prospective, randomized trial of autologous bone marrow transplantation and chemotherapy in multiple myeloma. *Intergroupe Francais du Myelome. N Engl J Med* 335: 91-97, 1996.
- 30 Bataille R, Jourdan M, Zhang XG, Klein B. Serum levels of interleukin-6, a potent myeloma cell growth factor, as a reflection of disease severity in plasma cell dyscrasias. *J Clin Invest* 84: 2008, 1989.
- Bataille R, Chappard D, Klein B. Mechanisms of bone lesions in multiple myeloma.
- 35 *Hem Onc Clin NA* 6: 285-295, 1992.

5

Bataille R, Barlogie B, Lu ZY, Rossi JF, Lavabre-Bertrand T, Beck T, Wijdenes J, Brochier J, Klein B. Biologic effects of anti-interleukin-6 murine monoclonal antibody in advanced multiple myeloma. *Blood* 86: 685-691, 1995.

10

Boyce BF, Yates AJP, Mundy GR. Bolus injections of recombinant human interleukin-1 cause transient hypocalcemia in normal mice. *Endocrinology* 125: 2780-2783, 1989.

15

Chauhan D, Uchiyama H, Urashima M, Yamamoto K, Anderson KC. Regulation of interleukin-6 in multiple myeloma and bone marrow stromal cells. *Stem Cells* 13: 35-39, 1995.

Epstein J. Myeloma phenotype: Clues to disease origin and manifestation. *Hem Onc Clin NA* 6: 249-256, 1992.

20

Garrett IR, Dallas S, Radl J, Mundy GR. A murine model of human myeloma bone disease. *Bone* 20: 515-520, 1997.

25

Gossler U, Jonas P, Luz A, Lifka A, Naor D, Hamann A, Holzmann B. Predominant role of alpha 4 integrins for distinct steps of lymphoma metastasis. *Proc. Natl. Acad. Sci. USA*. 93: 4821-4826, 1996.

30

Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93: 165-176, 1998.

Lobb R, Chi-Rosso G, Leone D, Rosa M, Newman B, Luhowskyj S, Osborn L, Schiffer S, Benjamin C, Dougas I, Hession C, Chow P. Expression and functional

- 5 characterisation of a soluble form of vascular cell adhesion molecule 1. *Biochem. Biophys. Res. Commun.* 178: 1498-1504, 1991.
- Lobb, R. and Hemler, M. The Pathophysiologic Role of alpha4 Integrins In Vivo. *J. Clin. Invest.*, 94: 1722-1728 (1994).
- 10 MacDonald BR, Mundy GR, Clark S, Wang EA, Kuehl TJ, Stanley ER, Roodman GD. Effects of human recombinant CSF-GM and highly purified CSF-1 on the formation of multinucleated cells with osteoclast characteristics in long-term bone marrow cultures. *J Bone Min Res* 1: 227-233, 1986.
- 15 Mbalaviele G, Chen H, Boyce BF, Mundy GR, Yoneda T: The role of cadherin in the generation of multinucleated osteoclasts from mononuclear precursors in murine marrow. *J Clin Invest* 95: 2757-2765, 1995.
- 20 Matsuura N, Puzon-McLaughlin W, Irie A, Morikawa Y, Kakudo K, Takada Y. Induction of experimental bone metastasis in mice by transfection of integrin alpha 4 beta 1 into tumor cells. *Am J Pathol* 148: 55-61, 1996.
- 25 Matsuzaki K, Udagawa N, Takahashi N, Yamaguchi K, Yasuda H, Shima N, Morinaga T, Toyama Y, Yabe Y, Higashio K, Suda T. Osteoclast differentiation factor (ODF) induces osteoclast-like cell formation in human peripheral blood mononuclear cell cultures. *Biochem Biophys Res Commun* 246: 199-204, 1998.
- 30 Mundy GR, Bertolini DR. Bone destruction and hypercalcemia in plasma cell myeloma. *Seminars Oncol* 3: 291, 1986.
- Mundy GR. Myeloma bone disease. *Eur. J. Cancer* 34: 246-251, 1998.

- 5 Papayannopoulou T, Nakamoto B. Peripheralization of hemopoietic progenitors in primates treated with anti-VLA4 integrin. *Proc. Natl. Acad. Sci. USA* 90: 9374-9378, 1993.
- 10 Qian F, Vaux DL, Weissman IL. Expression of the integrin $\alpha 4 \beta 1$ on melanoma cells can inhibit the invasive stage of metastasis formation. *Cell*, 77: 335-347, 1994. —
- Radl J, Croese JW, Zurcher C, van den Enden-Vieveen MM, de Leuw AM. Animal model of human disease. *Am. J. Pathol.* 132: 593-597, 1988.
- 15 Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Luthy R, Nguyen HQ, Wooden S, Bennett L, Boone T, Shimamoto G, DeRose M, Elliott R, Colombero A, Tan HL, Trail G, Sullivan J, Davy E, Bucay N, Renshaw-Gegg L, Hughes TM, Hill D, Pattison W, Campbell P, Boyle WJ, et al. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* 309-319, 1997.
- 20 Takahashi N, Yamana H, Yoshiki S, Roodman GD, Mundy GR, Jones SH, Boyde A, Suda T: Osteoclast-like cell formation and its regulation by osteotropic hormones in mouse bone marrow cultures. *Endocrinology* 122: 1373-1382, 1988.
- 25 Vanderkerken K, De Raeve H, Goes E, Van Meirvenne S, Radl J, Van Riet 1, Thielemans K, Van Camp B. Organ involvement and phenotypic adhesion profile of 5T2 and 5T33 myeloma cells in the C57BL/KaLwRij mouse. *Brit J Cancer* 76: 451-460, 1997.
- 30 Vaughan T, Williams AJ, Pritchard K, Osbourn JK, Pope AR, Earnshaw JC, et al. Human antibodies with sub-nanomolar affinities isolated from a large non-immunized phage display library. *Nature Biotechnology*. 14: 309-314, 1996.
- 35 Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, Mochizuki S, Tomoyasu A, Yano K, Goto M, Murakami A, Tsuda E, Morinaga T, Higashio K, Udagawa N,

5 Takahashi N, Suda T. Osteoclast differentiation factor is a ligand for
osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to
TRANCE/RankL. Proc Natl Acad Sci USA 95: 3597-3602, 1998.

10 Yoneda T, Alsina MM, Garcia JL, Mundy GR: Differentiation of HL-60 Cells into cells
with the osteoclast phenotype. Endocrinology 129: 683-689, 1992. —

15 Yoneda T, Lowe C, Lee CH, Gutierrez G, Niewolna M, Williams P, Izbicka E, Uehara
Y, Mundy GR: Herbimycin A, a pp60^{c-src} tyrosine kinase inhibitor, inhibits osteoclastic
bone resorption in vitro and hypercalcemia in vivo. J Clin Invest 91: 2791-2795, 1993.

5 7) CLAIMS:

1. A method for the treating multiple myeloma comprising administering to an individual a therapeutically effective amount of a composition comprising an antagonist of an interaction between an alpha4 subunit-bearing integrin and a ligand for an alpha4 subunit-bearing integrin.

2. The method of claim 1, wherein the antagonist is an alpha 4 integrin binding agent.

3. The method of claim 1, wherein the antagonist is an alpha4 integrin ligand binding agent.

15

4. The method of claim 2, wherein the alpha 4 integrin binding agent is selected from the group consisting of; a) an antibody homolog that antagonizes the interaction of both VLA-4 and alpha4 beta 7 with their respective alpha4 ligands; b) an antibody homolog that antagonizes the interaction of VLA-4 with its alpha4 ligand; and c) an antibody homolog that antagonizes

20 the interaction of alpha4beta7 with its alpha4 ligand.

5. The method of claim 4, wherein the antibody homolog is selected from the group consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments thereof.

25

6. The method of claim 3, wherein the alpha4 integrin ligand binding agent is an anti-VCAM-1 antibody homolog.

7. The method of claim 6, wherein the antibody homolog is selected from the group
30 consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments
thereof.

8. The method of claim 1, wherein the antagonist is a small molecule.

35 9. A method of claim 1, wherein the composition is administered at a

5 dosage so as to provide from about 0.1 to about 20 mg/kg body weight.

10. A method for inhibiting bone resorption associated with tumors of bone marrow, the method comprising administering to a mammal with said tumors an antagonist of an interaction between an alpha4 subunit-bearing integrin and a ligand for an alpha4 subunit-bearing integrin, in an amount effective to provide inhibition of said bone resorption.

11. The method of claim 10, wherein the antagonist is an alpha4 integrin binding agent.

12. The method of claim 10, wherein the antagonist is an alpha4 integrin ligand binding agent.

13. The method of claim 11, wherein the alpha4 integrin binding agent is an anti-VLA4 antibody homolog or anti-alpha4beta 7 antibody homolog.

14. The method of claim 13, wherein the antibody homolog is selected from the group consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments thereof.

15. The method of claim 12, wherein the alpha4 integrin ligand binding agent is an anti-VCAM-1 antibody homolog.

16. The method of claim 15, wherein the antibody homolog is selected from the group consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments thereof.

17. The method of claim 10, wherein the antagonist is a small molecule.

18. A method of claim 10, wherein the antagonist is administered at a dosage so as to provide from about 0.1 to about 20 mg/kg, based on the weight of the individual.

5

19. The method of claim 17, wherein the antagonist is administered in an amount effective to provide a dosage of small molecule of about 0.1 -30 mg/kg body weight.

20. A method of treating a subject having a disorder characterized by the presence of osteoclastogenesis, the method comprising administering to the subject an antagonist of an
10 interaction between an alpha4 subunit-bearing integrin and a ligand for an alpha4 subunit-bearing integrin, in an amount sufficient to suppress the osteoclastogenesis.

21. The method of claim 20, wherein the antagonist is an alpha4 integrin binding agent.
15

22. The method of claim 20, wherein the antagonist is an alpha4 integrin ligand binding agent.

23. The method of claim 21, wherein the alpha4 integrin binding agent is an anti-VLA4
20 antibody homolog or an anti-alpha4 beta 7 binding agent.

24. The method of claim 23, wherein the antibody homolog is selected from the group consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments thereof.
25

25. The method of claim 22, wherein the alpha4 integrin ligand binding agent is an anti-VCAM-1 antibody homolog.

26. The method of claim 25, wherein the antibody homolog is selected from the group consisting of a human antibody, a chimeric antibody, a humanized antibody and fragments thereof.
30

27. The method of claim 20, wherein the antagonist is a small molecule.

35 28. The method of claim 20, wherein the antagonist is administered at a

5 dosage so as to provide from about 0.1 to about 20 mg/kg body weight.

29. The method of claim 27, wherein the antagonist is administered in an amount effective to provide a dosage of small molecule of about 0.1 -20 mg/kg body weight.

10

1/6




FIG. 1

2/6





FIG. 2



4/6




FIG. 4

5/6




FIG. 5

6/6




FIG. 6